

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
1 July 2004 (01.07.2004)

PCT

(10) International Publication Number
WO 2004/054571 A1

(51) International Patent Classification⁷: **A61K 31/35**,
31/135, 9/52, 9/20, 47/36

(74) Agent: WANTE, Dirk; Johnson & Johnson NCB, Eu-
ropean Patent Group, Lenneke Marelaan 6, B-1932 Sint-
Stevens-Woluwe (BE).

(21) International Application Number:
PCT/EP2003/014474

(22) International Filing Date:
12 December 2003 (12.12.2003)

(25) Filing Language: English

(26) Publication Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO,
CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB,
GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG,
KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG,
MK, MN, MW, MX, MZ, NO, NZ, OM, PG, PH, PL, PT,
RO, RU, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT,
TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(30) Priority Data:
02080325.0 13 December 2002 (13.12.2002) EP
03075123.4 10 January 2003 (10.01.2003) EP

(84) Designated States (*regional*): ARIPO patent (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): CILAG
AG [CH/CH]; Hochstrasse 207, CH-8205 Schaffhausen
(CH).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): BACHMANN,
Dieter [DE/DE]; Buchenstrasse 87, CH-8212 Neuhausen
(CH). EIVASKHANI, Reza [DE/DE]; Wildalstrasse 71,
79108 Freiburg i.br. (DE). BRAUN, Christian [DE/DE];
Im Mühlezelge 36, 78239 Rielasingen-Worbingen (DE).
SPYCHER, René [CH/CH]; Moerlengasse 4, CH-8463
Benken (CH). STRONG, Brian [GB/GB]; Ledergasse 16,
CH-8232 Merishausen (CH).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: CONTROLLED RELEASE PREPARATIONS COMPRISING TRAMADOL AND TOPIRAMATE

(57) Abstract: This invention relates to an oral pharmaceutical preparation, suitable for dosing every 24 hours, comprising a sub-
strate, which substrate comprises a pharmaceutically effective amount of tramadol or a salt thereof and a pharmaceutically effective
amount of topiramate and wherein said substrate may be coated with a controlled release coating; said preparation having a specific
dissolution rate in vitro.

WO 2004/054571 A1

Controlled Release Preparations Comprising Tramadol and Topiramate

5

Field of the invention

This invention relates to an oral pharmaceutical preparation, suitable for dosing every 24 hours, comprising a substrate, which substrate comprises a pharmaceutically effective amount of tramadol or a salt thereof and a pharmaceutically effective amount of topiramate and wherein said substrate may be coated with a controlled release coating; said preparation having a specific dissolution rate in vitro.

15

Background of the Invention

A number of effective anticonvulsants including the compound 2,3:4,5-bis-O-(1-methylethylidene)- β -D-fructopyranose sulfamate, also known as topiramate, have been disclosed in U.S. Pat. No. 4,513,006. Topiramate is useful in the treatment of human epilepsy in that it is effective as adjunctive therapy or as monotherapy in treating simple and complex partial seizures and secondarily generalized seizures. Topiramate is currently marketed for the treatment of simple and complex partial seizure epilepsy with or without secondary generalized seizures.

Recent preclinical studies on topiramate have revealed previously unrecognised pharmacological properties, which suggest that topiramate is effective in treating some other disorders. One of these is neuropathic pain, which remains one of the "frontiers" of pain management. There is a significant unmet need for efficacious and tolerable pharmacotherapy, making neuropathic pain an area of intense research interest. The term "neuropathic pain" is applied to any acute or chronic pain syndrome in which the sustaining mechanism for the pain is believed to involve abnormal transmission (peripheral) or processing (central) of somatosensory input.

U.S. Pat. No. 5,760,007 discloses topiramate as useful for the treatment of neuropathic pain. US 5,935,933 discloses anticonvulsant derivatives useful in treating neuropathic pain, including but not limited to neuralgia.

- 5 A class of analgesic cycloalkanol-substituted phenol esters having a basic amine group in the cycloalkyl ring, are disclosed in U.S. Pat. No. 3,652,589. Among these is the compound (1R,2R or 1S,2S)-2-[(dimethylamino)methyl]-1-(3-methoxyphenyl) cyclohexanol, commonly known as tramadol, which is specifically disclosed therein.
- 10 A series of articles pertaining to the pharmacology, toxicology and clinical studies of tramadol are found in *Arzneim. Forsch.*, (Drug Res.), 1978, 28(1), 114. The Abstracts of the VIIth World Congress on Pain, April 1-6 (1990), disclose that tramadol hydrochloride is an orally active pure agonist opioid analgesic. However, clinical experience indicates that tramadol lacks many of the typical side effects of opioid
- 15 agonists, e.g., respiratory depression, constipation, tolerance and abuse liability. Tramadol's opioid activity without the typical side effects thereof makes it a very unique drug. Tramadol is currently marketed as an analgesic.

As an analgesic, tramadol has been combined with both opioid and non-opioid

20 analgesic drugs. Such compositions have exhibited synergistic effects in treating pain while using less of each ingredient to produce an equivalent degree of analgesia. Specifically, U.S. Pat. No. 5,516,803 discloses the composition of tramadol and a NSAID, particularly ibuprofen. U.S. Pat. No. 5,468,744 discloses tramadol plus any of oxycodone, codeine or hydrocodone and U.S. Pat. No. 5,336,691 discloses tramadol in

25 combination with acetaminophen.

WO-01/13904 relates to a pharmaceutical composition comprising a combination of a tramadol material and an anticonvulsant drug, in particular topiramate, and to the pharmacological use of the composition in treating conditions of pain and neurologic or

30 psychiatric disorders. The composition has improved properties, requiring less of each ingredient and producing a synergistic effect. EP-A-699 436 discloses controlled release preparations for oral administration containing tramadol or a salt thereof with a

specific in vitro dissolution profile. EP-A-914 823 discloses sustained release preparations with tramadol hydrochloride as active ingredient with a fatty alcohol as matrix forming agent. US-5,427,799 concerns sustained release compositions comprising an active ingredient, which may be an analgesic, in mixture with xanthan gum.

In animal studies using the Chung model of post-nerve constriction injury, both tramadol and topiramate are significantly active and reach 100% MPE as the dose is escalated. When topiramate and tramadol are co-administered in this model the ED₅₀ of both drugs is dramatically reduced, suggesting synergy of analgesic effect. The degree of synergy varies across ratios in this model with those ratios in which tramadol predominates displaying the greatest synergy.

In order to be effective in the treatment of neuropathic pain, it is necessary that sufficient amounts of both tramadol and topiramate are present at the receptor sites, in particular in the concentrations and concentration ratios required to produce the synergistic effect described in WO-01/13904. A particularly effective w/w ratio of both ingredients is in the range of about 1 : 1.5 to about 1 : 5, in particular from about 1 : 3 to about 1 : 5 of topiramate : tramadol. This w/w ratio should be reflected in the blood plasma levels of both these ingredients. Maintaining said blood plasma levels within this concentration ratio is a desirable goal to achieve. This goal becomes even more challenging when sustained release formulations are desired for administration every 12 or 24 hours, this because of the different half-life values of both agents. Indeed, the metabolism rate of topiramate is a relatively slow while that of tramadol is relatively fast. Thus, without special measures, the blood plasma level of topiramate will be largely in excess to that of tramadol so that the latter no longer is present in a sufficient amount to be effective and the synergistic effect no longer is present.

Compositions including combinations of topiramate and tramadol have been found to be relatively unstable, in particular due to the instability of topiramate. There is therefore a need to provide formulations containing both topiramate and tramadol, with increased stability. Moreover there is a need to provide a composition that releases both

topiramate and tramadol in a manner such that the blood plasma levels are sufficient for both active ingredients to be effective and to act synergistically. Providing such compositions is an object of this invention. A further object of this invention is to provide a method for treating conditions of pain, in particular neuropathic pain in
5 mammals.

Summary of the invention

10 This invention relates to a controlled release oral pharmaceutical preparation suitable for dosing every 24 hours comprising a substrate comprising a pharmaceutically effective amount of tramadol or a salt thereof and a pharmaceutically effective amount of topiramate; said preparation having a dissolution rate in vitro when measured using the Ph. Eur. Paddle Method at 75 rpm in 900 ml 0.05 M phosphate buffer at 37° C and
15 using high performance liquid chromatography (HPLC) :
between 15 and 35% tramadol released after 1 hour;
between 28 and 48% tramadol released after 2 hours;
between 47 and 67% tramadol released after 4 hours;
between 68 and 88% tramadol released after 8 hours;
20 between 79 and 99% tramadol released after 12 hours;
between 86 and 105% tramadol released after 18 hours;
about 100% tramadol released after 24 hours; by weight,
said preparation providing a therapeutic effect for about 24 hours after oral
administration.

25 In a preferred aspect there is provided an oral pharmaceutical preparation as defined above, said preparation having a dissolution rate as follows :
between 20 and 30% tramadol released after 1 hour;
between 33 and 43% tramadol released after 2 hours;
30 between 52 and 62% tramadol released after 4 hours;
between 73 and 83% tramadol released after 8 hours;
between 84 and 94% tramadol released after 12 hours;

between 91 and 100% tramadol released after 18 hours;
about 100% tramadol released after 24 hours; by weight.

In a most preferred aspect there is provided an oral pharmaceutical preparation as
5 defined above, said preparation having a dissolution rate as follows :

about 25% tramadol released after 1 hour;
about 38% tramadol released after 2 hours;
about 57% tramadol released after 4 hours;
about 78% tramadol released after 8 hours;
10 about 89% tramadol released after 12 hours;
about 96% tramadol released after 18 hours;
about 100% tramadol released after 24 hours; by weight.

In preferred embodiments the tramadol salt is tramadol hydrochloride.
15

In further embodiments the said substrate is a suitable matrix material in which
tramadol or its salt form is incorporated, said matrix material preferably comprising
xanthan gum.

20 In a further aspect, the invention concerns pharmaceutical preparations as described
herein comprising two or more phases. In certain embodiments, the major part of the
tramadol or its salt form and of the topiramate are in different phases of the said
pharmaceutical preparations. In these embodiments at least one phase may contain
either the major part of topiramate or the major part of tramadol or a salt-form thereof.

25 In particular embodiments, one phase contains the major part of tramadol or a salt
thereof and another phase contains the major part of topiramate. Further particular
embodiments are pharmaceutical preparations that take the form of a bi-phasic tablet
having a phase that comprises the major part of topiramate and another phase that
comprises the major part of tramadol or a salt form thereof. The phases in these
30 embodiments may take the form of layers.

In a specific aspect the invention concerns pharmaceutical preparations as described herein comprising two or more phases, wherein the tramadol or its salt form and the topiramate are in different phases of the said pharmaceutical preparations and none of the phases contains as well topiramate as tramadol or a salt-form thereof. In particular
5 embodiments, one phase contains the tramadol or a salt thereof and another phase contains the topiramate. Further particular embodiments are pharmaceutical preparations that take the form of a bi-phasic tablet having a phase that comprises the topiramate active ingredient and another phase that comprises the tramadol active ingredient or a salt form thereof.

10

In a particular aspect this invention concerns pharmaceutical preparations as defined herein, wherein said preparations are bi- or multi-layer tablets and wherein none of the layers contains as well topiramate as tramadol or a salt-form thereof. In particular
15 embodiments, the previously mentioned pharmaceutical preparations take the form of a bilayer tablet having a phase that comprises the topiramate active ingredient and another phase that comprises the tramadol active ingredient or a salt form thereof.

In a further aspect, there is provided a bi- or multiphasic tablet according to the invention containing an effective amount of topiramate having at least one phase
20 or layer that contains from about 20% to about 100%, in particular from about 30% to about 90% or from about 50% to 80% of polymeric matrix material. The latter preferably is xanthan gum.

In a particular embodiment, the tablets according to the present invention are coated
25 with an appropriate coating. The coating may be for taste masking or for other purposes.

The invention also provides preparations as defined herein which are capsules or sachets. The topiramate and/or the tramadol containing phase or phases in these
30 embodiments may take the form of pellets.

In another aspect the invention concerns a process for manufacturing the oral pharmaceutical preparation described herein comprising mixing tramadol hydrochloride, being incorporated in a suitable controlled release substrate, and topiramate, preferably formulated in a suitable carrier material solid form.

5

In a further aspect there is provided a process for manufacturing a bi- or multiphasic tablet in accordance with the invention, comprising compressing two or more pre-shaped phases in an appropriate compressing apparatus.

10 In a further aspect there is provided a process for manufacturing a bi- or multilayer tablet in accordance with the invention comprising compressing a suitable topiramate containing composition as to form a layer, laying tramadol containing matrix material on this topiramate containing layer, compressing the whole, and if desired laying further compositions of topiramate and/or further tramadol containing matrix material
15 thereon and each time subjecting the whole to a compression and if further desired coating the thus prepared dosage form.

In a further aspect there is provided a process for manufacturing a bi- or multilayer tablet in accordance with the invention comprising compressing tramadol containing
20 matrix material as to form a layer, laying a suitable topiramate containing mixture on this tramadol containing matrix material layer, compressing the whole, and if desired laying further compositions of topiramate and/or further tramadol matrix material thereon and each time subjecting the whole to a compression and if further desired coating the thus prepared dosage form.

25

Furthermore, the invention concerns a method of treating a warm blooded animal suffering from neuropathic pain, said method comprising the administration of a pharmaceutical preparation as described herein.

30

Detailed description of the invention

As mentioned herein, any % is weight-by-weight, relative to the total weight of the preparation or the formulation.

5

Tramadol is the compound (1R,2R or 1S,2S)-2-[(dimethylamino)methyl]-1-(3-methoxyphenyl)-cyclohexanol. Preferably tramadol is used as a salt form, in particular its hydrochloride salt. Tramadol is commercially available from Gruenenthal or may be made by the process described in U. S. Patent No. 3,652,589, which is herein
10 incorporated by reference.

15

Topiramate is the compound 2,3,4,5-bis-O-(1-methylethylidene)- β -D-fructopyranose sulfamate and can be prepared according to processes described in U.S. Pat. No. 4,513,006.

20

The controlled release preparation should release tramadol *in vitro* in the quantities outlined above. These are obtained by measurement using the Ph. Eur. Paddle Method at 75 rpm in 900 ml 0.05 M phosphate buffer with a pH value of 6.8 (USP) at 37° C and using high performance liquid chromatography (HPLC). In the latter a suitable
20 detection system is used such as, for example, UV detection at an appropriate wavelength, e.g. at 270 nm in the case of tramadol, or with a refractive index detector. Alternatively, release of the active ingredients and in particular release of tramadol can be measured by using *in situ* measurement with a fiber optic dissolution system, using the second derivative correction method at a suitable wavelength range, which, in case
25 of tramadol is in the range of 283 to 289 nm.

30

The controlled release preparations according to the invention should be suitable for dosing every 24 hours. The 'term suitable for dosing every 24 hours' means that the dosage forms should be such that they can be administered every 24 hours and give
30 effective blood plasma concentrations of both active ingredients such that they are effective to treat neuropathic pain over a period of 24 hours. The controlled release

preparations according to the invention can be dosed every 24 hours but also can be dosed differently, e.g. every 12 hours (b.i.d.) or every 8 hours (t.i.d.).

As mentioned above, in order to act synergistically, the ratios of the blood plasma
5 levels of topiramate : tramadol should be within certain ranges and in particular the blood plasma levels of these ingredients should be in the range of about 1 : 1.5 to about 1 : 5, in particular from about 1 : 3 to about 1 : 5 of topiramate : tramadol. It has been found that an optimum ratio for these ingredients is about 1 : 3 topiramate : tramadol.

10 It further has been found that when tramadol is released *in vitro* in the quantities outlined above, upon multiple administrations during specific periods of time, e.g. every 12 hours, or, which is preferred, every 24 hours, the plasma concentrations of tramadol *in vivo* reach a steady state and are constant within certain ranges during an extended period of time. It has additionally been found that in pharmaceutical
15 preparations as described herein, containing tramadol or a salt thereof and topiramate, when the release of tramadol follows the release pattern as outlined above, the ratio of the plasma concentrations of topiramate vis à vis tramadol is constant within certain ranges. This ratio has been found to approximate 1 : 3 (w/w) or upon selection of the appropriate concentrations of the topiramate and tramadol, or its salt form, and of the
20 substrate and other carriers in the pharmaceutical dosage form, this ratio may be 1 : 3 or about 1 : 3. This equally means that upon multiple administrations of the pharmaceutical preparations of this invention, also during specific periods of time, e.g. every 12 hours, or, which is preferred, every 24 hours, the plasma concentrations of topiramate *in vivo* reach a steady state and are constant within certain ranges during a
25 long period of time. This will allow both agents to act synergistically and therefore to be effective in combating neuropathic pain during a longer period of e.g. 12 or even 24 hours upon administration of the pharmaceutical preparations in accordance with the invention.

30 As used herein, 'constant within certain ranges' means that there can be small fluctuations of the plasma concentrations or of the ratio of the plasma concentrations within an acceptable range, e.g. within 30 % in particular, within 20 %, further in

particular within 10 %. This alternatively can be expressed by the 'fluctuation index' which is defined as follows :

$$5 \quad F_i = \frac{C_{\max} - C_{\min}}{C_{Av}}$$

wherein F_i is the fluctuation index, C_{\max} the maximal plasma concentration, C_{\min} the
10 minimal plasma concentration, and C_{Av} the average plasma concentration. The fluctuation index can vary but for example is 1.3, or 1.2 or even 1.1.

The *in vitro* release quantities of tramadol outlined above are based on an analysis of the *in vivo* plasma concentration data of topiramate and on the *in vitro* / *in vivo*
15 correlation of plasma concentrations and *in vitro* release quantities of tramadol. Administration of an effective amount of topiramate will show a particular course of the plasma concentrations. Ideally, tramadol should follow the same course so that the ratio of the plasma concentrations of both agents remains more or less constant. By measuring the correlation of plasma concentrations and *in vitro* release quantities of
20 tramadol it has been found that it is possible to predict which plasma concentration correlates with a given *in vitro* release quantity and vice versa (this method can be referred to as reversed IVIVC, i.e. reversed *in vitro* *in vivo* correlation). This allows a reverse calculation of the *in vitro* release quantities that would cause *in vivo* plasma concentrations, which run in parallel with the plasma concentrations of topiramate.

25 This invention relates to controlled release oral pharmaceutical preparations as specified herein comprising a substrate comprising a pharmaceutically effective amount of tramadol or a salt thereof and a pharmaceutically effective amount of topiramate. As used herein, the term 'substrate' refers to any material or combination of materials, or
30 forms thereof, that results in the specified release pattern of tramadol.

Typically, the pharmaceutical preparations according to the present invention comprise tramadol or its salt form in a suitable controlled release form, which may be any form that affords release of tramadol within the ranges specified above.

In certain embodiments, the preparations of the invention comprise controlled release forms wherein the tramadol or its salt form is incorporated in a suitable matrix, which may be a controlled release matrix or a normal release matrix having a controlled
5 release coating. The controlled release form may take various forms, e.g. particles of different sizes, pellets (or beads), tablets, phases within a larger unit such as layers or sections of other shape within a larger unit (e.g. as in a multi-layer or a bull-eye tablet). A number of such formats as well as the unit dosage forms in which these can be incorporated will be outlined in more detail hereinafter.

10

As used herein the term 'phase' refers to a defined three dimensionally shaped section in a tablet dosage form that contains the same material and wherein each phase is separated from the other. Examples of phases are layers, which are incorporated in bi- or multi-layer tablets. Other examples are cylindrical, spherical or other
15 tridimensionally shaped sections that can be present in tablets. This gives rise to different tablet formats such as the so-called 'bull-eye' tablets, or concentric tablets (a central cylindrically shaped section completely surrounded with one or more further cylindric layers (i.e. a ring-like combination), or 'coated' tablets wherein the coating is a layer completely surrounding a tablet nucleus and the like tablet formats. Preference
20 is given to bi- or multi-layer tablets.

Preferred embodiments are tablets that contain at least two phases, in particular tablets that contain at least two layers.

25 In particular embodiments, the major part of the tramadol or its salt form and of the topiramate are in different phases of the said pharmaceutical preparations. In said embodiments a at least one phase may contain either the major part of topiramate or the major part of tramadol or a salt-form thereof. In particular embodiments, one phase contains the major part of tramadol or a salt thereof and another phase contains the
30 major part of topiramate. As used herein, 'major part' means that the major quantity of the tramadol or its salt form or of topiramate is present in a particular phase. Preferably the term 'major part' refers to a situation where at least more than about 90 % of the

concerned active ingredient is present in a particular phase, for example more than 95%, or more than 98%, or more than 99%, or even more than 99.5%. The same applies to the situation take particular forms such as layers.

- 5 Most preferably a phase containing one of both active ingredients should contain only a minute amount of the other active ingredient, or even none of the other active ingredient, for example a phase may contain topiramate and a minute amount, e.g. less than 1%, or less than 0.5% of tramadol or a salt form thereof, or vice versa.
- 10 Preferably a phase comprising tramadol or a salt form thereof is adjacent to a phase containing topiramate.

Of particular interest are tablets that are biphasic, the latter being preferred, or multiphasic, e.g. having 3, 4, 5 or more phases. At least one layer should comprise
15 tramadol or a salt form thereof but in case of multiphasic tablets, more than one layer comprising tramadol or a salt form can be present. Of still further interest are those preparations in which one or more of the phases are layers.

Particularly preferred embodiments are tablets wherein topiramate is present in
20 amounts from about 10 mg to 500 mg topiramate per unit, preferably from about 25 mg to about 200 mg of topiramate per unit, e.g. tablets having 25, 50, 100 or 200 mg per unit.

In a particular aspect, the tablets of the invention contain an effective amount of
25 topiramate, wherein the tablets have at least one layer that contains from about 20% to about 100%, in particular from about 30% to about 90% or from about 50% to 80% of polymeric matrix material. The hygroscopic matrix material containing layer may contain other ingredients such as the ingredients mentioned hereinafter.

Controlled release matrix

Where the matrix is a controlled release it may comprise suitable digestible hydrophilic or hydrophobic polymeric or non-polymeric materials.

5

Examples of such polymeric materials are hydrophilic or hydrophobic polymers, such as polysaccharides, in particular gums (further in particular pH dependent gums), cellulose ethers, especially alkylcelluloses, in particular C₁-C₆ alkyl cellulose, especially ethyl cellulose, acrylic resins, protein-derived materials, polyalkylene glycols, and the like. Preferred are the polysaccharide gums such as xanthan gum.

Examples of non-polymeric materials that can be used are digestible lipids having a long chain alkyl moiety, which may be straight or branched, saturated or unsaturated, substituted or unsubstituted. Of particular interest are C₈₋₅₀, especially C₁₂₋₄₀ lipids.

15 Examples comprise fatty acids, fatty alcohols, glyceryl esters of fatty acids, mineral and vegetable oils and waxes. Lipids having a melting point of between 25 and 90°C are preferred.

The tramadol controlled release form may conveniently contain between 1% and 90%, in particular from 10% to 90%, further in particular from 20% to 80% (by weight) of one or more hydrophilic or hydrophobic polymers or digestible lipids. In the embodiments where the polymeric material is a polysaccharide gum such as xanthan gum, alginate or gum Arabic, the preparation may contain between 20% and 90%, in particular from 30% to 80%, (by weight) of xanthan gum. As used herein, 'alginate'

25 refers to alginate or its salts, in particular to its alkali metal salts such as sodium or potassium salts. In the embodiments containing polyalkylene glycols, the preparation may in particular contain up to 60% (by weight) of one or more polyalkylene glycols. In further particular embodiments, the preparation may contain up to 60% (by weight) of at least one digestible, long chain lipid.

30

Of particular interest are controlled release matrixes comprising xanthan gum optionally in mixture with other gums, in particular with other pH dependent gums such as, for example, alginate.

Optionally, the controlled release matrix may also contain other pharmaceutically acceptable ingredients which are conventional in the pharmaceutical art such as diluents (in particular lactose), lubricants, binders, granulating aids, colorants, 5 flavorants, surfactants, pH adjusters, anti-adherents and glidants (e.g. colloidal silica), and plasticizers (e.g. dibutyl sebacate) and other suitable ingredients (e.g. ammonium hydroxide, oleic acid).

The tramadol controlled release form may conveniently be film coated using any film 10 coating material conventional in the pharmaceutical art. A film coat is added e.g. as a finish, for coloring purposes or taste masking or a combination of these. Preferably an aqueous film coating is used.

Alternatively, the tramadol controlled release form may comprise a normal release 15 matrix, as a core, and further having a controlled release coating. In such embodiments the tramadol form may be prepared via art known procedures, e.g. by a suitable granulation process followed by compression, or by direct compression, followed by a coating step with a coating material that ensures controlled release. In such preparations, the tramadol normal release section or core of the preparation may 20 contain any of the usual ingredients usually employed to make such normal release sections or cores. Any of the ingredients mentioned hereinafter with respect to the incorporation of topiramate in the preparations of the invention can be conveniently employed.

25 The release profile of tramadol can be adjusted in a number of ways. For instance a higher loading of the drug will be associated with increased initial release rates. By selecting particular ingredients and by controlling the relative amounts thereof in the preparation it is possible to adjust the release profile of the tramadol. Such particular ingredients for example are the matrix materials mentioned above, e.g. the polymeric 30 materials mentioned above.

Pellets

In further embodiments, the tramadol controlled release form comprises spherical pellets containing the active ingredient and a spheronizing agent. The pellets may be film-coated or not. The spheronizing agent may be any suitable pharmaceutically acceptable material, which can be spheronized together with the active ingredient to form pellets. The term 'spherical pellet' is meant to comprise pellets, beads or spheroids that are more or less of regular shape. In particular embodiments of the invention the shape is round or about round, i.e. having or approaching the shape of a small sphere.

The average size of the pellets may vary but preferably the diameter is in the range of about 0.1 mm to 3 mm, in particular from about 0.5 mm to about 2 mm, more preferably about 1 mm.

15

The size distribution of the pellets may vary but in general it is preferred that it has limited variation. It may vary between within a range of 10 to 20%. The size distribution may vary in a statistical manner, i.e. in a bell-shaped curve wherein e.g. 90% or e.g. 95% of the number of pellets are within a size range that varies between about 10% to about 20% of the average sizes mentioned above.

The tramadol or its salt form is present in an amount, which is in the range of from about 0.1 to about 50%, in particular from about 1 to about 40%, more in particular from about 10 to about 35%, w/w relative to the total weight of the pellet.

25

The pellets of the invention may further comprise an appropriate carrier which may be any carrier known in the art used for making pellets. Particular carrier materials are spheronizing agents that may be any suitable pharmaceutically acceptable material, which may be spheronized together with the active ingredient to form pellets. A preferred spheronizing agent is microcrystalline cellulose. The microcrystalline cellulose used may suitably be, for example, the product sold under the tradename 'AvicelTM'. The spheronizing agent is present in an amount, which is in the range of

30

from about 25% to about 90%, in particular from about 35% to about 70% w/w, relative to the total weight of the pellet.

Optionally the pellets may contain other pharmaceutically acceptable ingredients such as binders, bulking agents and colorants. Suitable binders, some of which may also contribute to the controlled release properties of the pellets, include water-soluble polymers, e.g. water-soluble hydroxyalkyl celluloses such as hydroxypropyl cellulose, or water insoluble polymers, such as acrylic polymers or copolymers, or alkyl celluloses such as, for example, ethylcellulose. Suitable bulking agents include lactose or colloidal silicon dioxide. The amount of these other ingredients in the pellets will be relatively small, e.g. lower than 30%, or 20%, or even lower than 10% or 5% w/w relative to the total weight of the pellet.

The pellets for use in the preparations of the present invention are made by an extrusion process followed by spheronization. The mixture used in the extrusion process comprises active ingredient, a suitable carrier material and other optional ingredients, and a suitable lubricant. The lubricant usually is water and the mixture for extrusion typically is converted into a granulate. After extrusion, the extrudate is spheronized to obtain pellets. If desired, the latter may be coated with a suitable coating material.

20

Tramadol is known to usually form a sticky mass upon contact with water and/or the other excipients used in the extrudate mixture. More in particular, tramadol seems to act as an additional binder in the mixture that is extruded and spheronized. This can be avoided by the addition of a dry lubricant. Apart from providing lubrication, the dry lubricant also allows the material to be extruded at a much lower moisture content thereby reducing the sticking observed in the spheronizer.

25

Further embodiments thus are spherical pellets for sustained release comprising tramadol or a salt thereof, a spheronizing agent and dry lubricant. In a further aspect, said pellets have a low water content. If desired, the pellets may be coated.

30

The dry lubricant in particular is a mono-, di- or triglyceride, or mixtures thereof. Suitable mono-, di- or triglycerides are the mono-, di- or triesters of glycerine and one or more fatty acids. The mono-, di- or triglycerides may contain the same or different fatty acid residues or mixtures thereof, e.g. technical mixtures obtained from
5 saponification of natural oils. Of particular interest are fatty acid triglycerides wherein the fatty acid residue has from 12 to 30 carbon atoms and is saturated or partially unsaturated or may be substituted, e.g. with one or more hydroxy functions. Preferred are mono-, di- or triglycerides derived from C₁₈₋₃₀ fatty acids, in particular derived from C₂₂₋₂₆ fatty acids. Of particularly preferred interest are behenic acid mono-, di- or
10 triglycerides.

The dry lubricant preferably is solid at room temperature and has a melting point or melting range which is in the range of 60 °C to 90 °C, in particular is in the range of 70 °C to 80 °C. A particularly suitable dry lubricant is the glyceride mixture sold under
15 the trade name 'CompritolTM 888ATO' which is a mixture of glyceryl mono-, di- and tribehenate, the dibehenate fraction being predominant, and having a melting range of about 69 – 74 °C.

Preferably, the dry lubricant is selected such that it does not impact the dissolution
20 behavior of the active ingredient.

The dry lubricant is present in an amount, which is in the range of from about 2% to about 50%, in particular between 10% and 35% w/w, relative to the total weight of the
25 pellet.

Of particular interest are pellets that have a low water content. In particular embodiments, the water contents in the pellets is lower than 5%, more in particular lower than 3%, w/w relative to the total weight of the pellet.

30 The tramadol spherical pellets, containing a dry lubricant, can be prepared by a process comprising extruding a mixture of the active ingredient with a suitable carrier in the presence of a dry lubricant and spheronizing the extrudate, wherein the dry lubricant is

a triglyceride. The amount of dry lubricant in this mixture may vary but in general is comprised between 10% and 35% (w/w). A small amount of water may be added to the mixture. In a particular execution, the amount of water is 5% or lower, or 3% or lower, or 1.5% or lower, w/w, relative to the total weight of the mixture for extrusion. In a
5 specific process the pellets are subsequently coated with a suitable coating.

The ingredients may be mixed together in any given sequence. In one embodiment, the dry lubricant is added to a mixture of active ingredient and the carrier material at room temperature. The mixture is subsequently extruded through a small orifice. The
10 diameter of the latter is in relation to the size of the pellets that are eventually produced from the extrudate. In one embodiment, the diameter of the orifices is in the range of 0.5 mm to 2.0 mm. The extrusion may be done at slightly elevated temperature but preferably is performed without applied heating. The extrudated material is subsequently placed into a spheronizer where it is spun at high speed.

15 In specific embodiments of the invention, the pellets (or spheroids), with or without dry lubricant, are subsequently coated with a suitable coating using art known methods. The coating can either be a functional coating or a diffusion controlling coating.

20 A functional coating may be applied for e.g. taste masking, protection of the pellets, to have improved stability (shelf-life) or for identification (for example by coloring). Functional coating often will be film coating, using any film coating material conventional in the pharmaceutical art. Preferably an aqueous film coating is used.

25 Diffusion controlling coatings are designed to achieve a target release profile such as controlled or sustained release permitting release of the active ingredient at a controlled rate in an aqueous medium. Suitable controlled or sustained release coating materials include water-insoluble waxes and polymers such as polymethacrylates, for example the EudragitTM polymers, or water insoluble celluloses, in particular alkyl celluloses
30 such as ethylcellulose. Optionally, water-soluble polymers such as polyvinylpyrrolidone or water-soluble celluloses such as hydroxypropylmethyl-cellulose or hydroxypropylcellulose may be included. Further components that may be

added are water-soluble agents such as polysorbate. Of particular interest is ethylcellulose (EC). Preferably, a suitable plasticizer is added. A coating material that is particularly suitable is the coating material sold under the trade name SureleaseTM (Colorcon), which is a dispersion of ethylcellulose.

5

Because of the bitter taste of the tramadol active ingredient, the pellets may be coated for taste-masking purposes although this may be of less importance if the pellets are used in a capsule dosage form.

10 Tramadol coated on beads

Alternatively the tramadol or its salt-form may be coated onto inert non-pareil beads, in particular onto sugar beads, and the drug loaded beads coated with a material, which permits control of the release of the active ingredient into the aqueous medium.

15

Topiramate

The topiramate in the preparations can be present throughout the pharmaceutical preparations of the invention or in particular sections thereof. In particular
20 embodiments it is present in one or more phases of the preparations. Preferably, the topiramate is present in one or more phases that do not contain tramadol.

The phases can take a variety of forms, e.g. sections in a tablet, or they can take the form of pellets. These forms can be prepared following art-known procedures. In the
25 particular case of sections in a tablet preparation, procedures can be applied such as granulation followed by partial or complete compression, or direct partial or complete compression.

Usually, the topiramate is formulated into a suitable formulation. This is prepared by
30 mixing topiramate with suitable ingredients into different formulation types such as powders, granulates, pellets and the like. The powder or granulate formulations may be compressed partially or completely to form appropriate phases for incorporation in bi-

or multi-phasic preparations. Particular phases are layers for incorporation in bi- or multi-layer tablets. Most preferably, the topiramate formulation will be for immediate release, i.e. the ingredients and the formulation form are selected such that release of topiramate is as quickly and as complete as possible. Ideally, release is 100 % after a
5 short period of time, e.g. within ½ hour.

In tablet preparations, suitable tableting excipients may be added e.g. one or more of the standard excipients such as diluents, lubricants, binding agents, flow aids, disintegrating agents, surface active agents or water soluble polymeric materials.
10 Suitable diluents are e.g. microcrystalline cellulose, lactose and dicalcium phosphate. Suitable lubricants are e.g. magnesium stearate and sodium stearyl fumarate. Suitable binding agents are e.g. hydroxypropyl methyl cellulose, polyvidone and methyl cellulose. Suitable disintegrating agents are starch, sodium starch glycolate, crospovidone and croscarmallose sodium. Suitable surface active are Poloxamer 188®,
15 Polysorbate 80 and sodium lauryl sulfate. Suitable flow aids are talc, colloidal anhydrous silica.

Double Layer Tablets

20 One particular execution of the controlled release preparations of the present invention are double layer (or bilayer) tablets. These comprise one layer containing tramadol dispersed in a suitable matrix and another layer that contains topiramate.

The topiramate containing layer preferably is composed of excipients typically used for
25 topiramate oral dosage forms such as tablets. Examples of such excipients comprise any of those mentioned above in relation to the formulation of topiramate.

The tramadol layer comprises any of the controlled release matrix materials described above. The matrix may in particular comprise a polysaccharide, more in particular a
30 gum, still more in particular xanthan gum. Or alternatively, the tramadol layer may consist essentially of polysaccharide, more in particular a gum, still more in particular xanthan gum.

The tramadol layer may contain from about 10 mg to 100 mg tramadol hydrochloride per unit, preferably from about 15 mg to about 75 mg of tramadol hydrochloride per unit, or from about 25 mg to about 65 mg of tramadol hydrochloride per unit. In case of application of tramadol-free base or other salts, an equivalent amount of active is used.

5

In particular embodiments of this invention, the tramadol layer contains an effective amount of tramadol, or a pharmaceutically acceptable salt thereof, dispersed in a matrix wherein the matrix contains from about 20% to about 90%, in particular from about 30% to about 80% of a polysaccharide, more in particular a gum, still more in particular xanthan gum. The percentages mentioned herein are w/w relative to the total weight of the dosage form.

10

The tramadol layer may additionally contain further ingredients such as the ingredients mentioned previously in relation to the topiramate layer, in particular starches, kaolin, lubricants, binders and the like. Preferred additional carriers are lubricants, e.g. magnesium stearate, flow enhancers or fillers, e.g. silica (silicon dioxide), fillers such as sugars, in particular lactose, titanium dioxide and the like.

15

In further particular embodiments of this invention, the tramadol layer contains as further ingredients lactose as a filler and magnesium stearate as a lubricant. Lactose is added to improve compressibility of the blend. Magnesium stearate is added to avoid tablet sticking on the lower or upper punch during the compression. The concentration of magnesium stearate in the tramadol layer may vary but good results are obtained when adding it in amounts ranging from about 0.5 to about 1.0% (w/w relative to the total weight of the dosage form). The concentration of lactose in the tramadol layer may vary but good results are obtained when adding it in amounts which range from about 5% to about 80%, preferably from about 10 % to about 65%, more preferably from about 20% to about 50% (w/w relative to the total weight of the dosage form).

20

25

The tramadol layer can be prepared by mixing tramadol or its salt form with the polysaccharide, more in particular the gum, still more in particular with xanthan gum while adding optional ingredients. The latter may also be added after the mixing of

30

tramadol and xanthan gum. The thus obtained mixtures are subsequently compressed, either by direct compression, which is preferred or by preparing a granulate and subsequent compression.

- 5 It has been found that when using tramadol and xanthan gum mixtures, the tablets can be prepared by direct compression. The mixtures for direct compression preferably contain a lubricant, in particular magnesium stearate. They may additionally contain a filler, in particular a sugar such as lactose. They may furthermore contain a flow enhancer such as colloidal silica (silicon dioxide). In the mixtures for direct
- 10 compression the lubricant preferably is present in concentrations in the range of about 0.75% to about 1.0%. The filler is present in concentrations from about 5% to about 80%, preferably from about 10% to about 65%, more preferably from about 20% to about 50%. The flow enhancer is present in concentrations from about 0.4% to about 0.6%, preferably about 0.45% to about 0.50%. All percentages herein are w/w relative
- 15 to the total weight of the tramadol containing phase or phases.

- Particular embodiments of the invention are coated tablets, in particular film-coated tablets. Coated tablets are easier to swallow than uncoated tablet cores, are usually easier to distinguish from other tablets - in particular when the film-coat contains a dye
- 20 or a pigment -, and may furthermore have an improved stability (shelf-life). In the present instance coating is mainly is for taste masking purposes because of the bitter taste of tramadol. Coatings are applied using art known methods using art known materials usually applied for this purpose. Particularly attractive coating products are based on suitable film-forming polymers such as hydroxypropylmethylcellulose
- 25 (HPMC) or polyvinylalcohol (PVA). Preferably, a plasticizer is added. Examples of suitable plasticizers are polyethylene glycol or derivatives thereof such as polyethoxylated alkylglycerides, e.g. polyethoxylated stearyl monoglyceride, in particular the material sold under the trade name MacroglTM. Further ingredients may be added to the coating such as fillers, dyes or pigments, flavors, sweeteners and the
- 30 like components. Examples of such further ingredients are lactose, titanium dioxide, starch and the like. Particularly suited as coating materials are the OpadryTM materials

which mainly contain the before mentioned materials and further ingredients such as plasticizers, e.g. polyethylene glycol.

In a preferred embodiment first the tramadol layer is produced by direct compression
5 whereupon topiramate granules are placed on top of the compressed tramadol layer as to form a second layer whereupon the whole is compressed to form a bi-layer tablet.

In particular embodiments there are provided bi-layer tablets comprising a tramadol layer and a topiramate layer wherein both layers are separated by a suitable layer that
10 may function as an isolator. This third layer may be comprised of suitable inert materials such as cellulose or lactose. Such embodiments may be prepared by first producing the tramadol layer by partial or complete compression of a suitable tramadol containing mixture whereupon the isolator material is put on the tramadol layer followed by a second compression, whereafter a suitable topiramate containing mixture
15 is put on top of the isolator layer as to form a third layer whereupon the whole is compressed to form a tri-layer tablet. The suitable tramadol containing mixture or suitable topiramate mixture may be a powder suitable for direct compression or a granulate obtained by a granulation process. The isolator layer may be desirable e.g. to avoid certain interactions between the components in each layer or to shield off
20 humidity.

Multi-layer tablets

Further embodiments are multi-layer tablets having multiple layers of topiramate and
25 tramadol, optionally separated by one or more isolator layers.

Further tablet formulations

In still further embodiments are tramadol tablets coated with a topiramate coating. In
30 this type of preparations, a suitable core containing tramadol or a salt thereof in a controlled release form is coated with a topiramate-containing coating, e.g. by spraying

with a suitable liquid formulation that contains topiramate. The core itself can be a tablet or another shaped phase.

5 Still further embodiments of the invention are so-called 'bull-eye' tablets, which are tablets with a cavity in which another tablet fits. The tablet with the cavity in particular is U-shaped. The tablet with a cavity may contain the tramadol and the other tablet the topiramate or vice versa. Bull-eye tablets can be made following art-known procedures using specially adapted punches in a tableting machine.

10 In any of the preparations that are tablets, the latter may be coated with a suitable coating material.

Preparations with pellets

15 Further embodiments of this invention are dosage forms comprising tramadol formulated in pellets, hereafter referred to as 'tramadol pellets'. The tramadol pellets may be prepared according to methods as described above and may be coated, if desired.

20 The tramadol pellets in turn may be coated with a topiramate containing coating, e.g. by spraying the tramadol pellets with an appropriate formulation containing topiramate. These tramadol pellets with a topiramate coating may be filled into capsules.

25 The tramadol pellets can be filled in capsules together with an appropriate formulation of topiramate e.g. formulated as a powder, granulate, or formulated itself as a pellet. The tramadol pellets and the topiramate formulation may be filled into the capsule in any give sequence, first the tramadol pellets followed by the topiramate formulation or vice versa or the two together or the two together as a mixture, e.g. a mixture of topiramate and tramadol pellets.

30

In further embodiments there are provided capsules containing tramadol pellets and one or more topiramate tablets. The topiramate tablets will evidently be of such size and

shape that it fits into a capsule. Preferably only one topiramate tablet is filled into one capsule.

5 In still further embodiments there is provided a so-called 'capsule into capsule' dosage form, i.e. a capsule containing a suitable topiramate formulation is put into a bigger capsule containing tramadol pellets. Or vice versa, a capsule containing tramadol pellets is put into a bigger capsule containing a suitable topiramate formulation. A suitable topiramate formulation can be a powder or a pellet formulation.

10 Still other embodiments are sachets filled with amounts of tramadol pellets and a suitable topiramate formulation.

In still another aspect, the invention concerns a process for manufacturing a pharmaceutical dosage form, said method comprising filling the tramadol pellets into a
15 suitable container and further adding a suitable topiramate formulation. In a preferred aspect the container is a capsule. Another type of container is a sachet.

In a particular embodiment, the invention provides unitary dosage forms, which comprise tramadol hydrochloride pellets as described herein in an amount that is such
20 that the dosage form contains an effective amount of tramadol hydrochloride. Particular embodiments of such dosage forms may contain from about 10 mg to 100 mg tramadol hydrochloride per unit, preferably from about 15 mg to about 75 mg of tramadol hydrochloride per unit, or from about 25 mg to about 65 mg of tramadol hydrochloride per unit.

25

The pharmaceutical preparations of the invention have a particular dissolution rate of the active ingredients in vitro and provide an effective therapeutic effect for a sufficiently long period of time, in particular such that the preparation is suitable for administering every 24 hours. Other dosing frequencies are also possible, e.g dosing
30 every 12 hours.

In a further aspect, the invention concerns a method for treating conditions of pain, in particular neuropathic pain, or a method of treating neurological or psychiatric disorders, in mammals in need of such treatment, said method comprising the administration to said mammals of a pharmaceutical preparation as described herein.

5

Preferably, the pharmaceutical preparations of the invention are suited for dosing every 24 hours.

Examples

10

Example 1

Double Layer Tablet Preparation:

Tramadol / Xanthan gum layer:

15

Actives and Excipients	mg/Tablet
Tramadol	45.00
Xanthan gum	160.00
Lactose (Fast Flo TM)	139.92
Magnesium Stearate	3.50
Silicon Dioxide	1.58
Total	350.00

Topiramate layer:

Active and Excipients	mg/Tablet
Topiramate	15.00
Lactose Monohydrate	18.51
Pregelatinized Starch	3.84
Microcrystalline Cellulose	8.25

Sodium Starch Glycolate	2.40
Magnesium Stearate	0.72
Lactose (Fast Flo™)	47.75
Total	96.46

The double layer tablet weight is: 446.46 mg.

Dry blend preparation prior to compression:

5

After blending the dispensed amount of tramadol HCl, xanthan gum and lactose for 20 minutes, the dispensed and sieved amount of magnesium stearate and silicium dioxide were added and blended for further 3 minutes. According to the topiramate strength needed in tablets the topiramate granules (with a drug load of 31.1% Topiramate) were diluted by adding lactose to the desired strength. The tramadol HCl dry blend and the topiramate blend were compressed to double layer tablets on a double layer rotary press tableting machine.

15 Example 2

Preparation of tramadol HCl pellets.

A dry blend of 1400 mg of tramadol hydrochloride, 1400 mg of microcrystalline cellulose and 1200 mg of glyceryl behenate (Compritol 888 ATO™, Gattefosse) is wet massed with approximately 60 mg of water and extruded through a small orifice (approx. 1 mm). The extruded material is placed into a spheroniser where it is spun at high speed (pellet speed of between 5 and 20 ms⁻¹). During this step the extrudate breaks and rounds into pellets, the size being determined by the size of the extrusion orifice. The pellets are coated uniformly with the coating material sold under the trade name Surelease™ (Colorcon), which is a dispersion of ethylcellulose.

25

Preparation of topiramate pellets

- A blend of 1400 mg of topiramate, 1400 mg of microcrystalline cellulose and 1260 ml is wet massed and extruded through a small orifice (approx. 1 mm). The extruded material is placed into a spheroniser where it is spun at high speed (pellet speed of between 5 and 20 ms^{-1}). During this step the extrudate breaks and rounds into pellets, the size being determined by the size of the extrusion orifice. The pellets are coated uniformly with the coating material sold under the trade name SureleaseTM (Colorcon).
- 3000 mg of the tramadol HCl pellets and 1000 mg of the topiramate pellets, both as prepared above are mixed. The thus prepared spherical pellet mixture is filled into capsules, at 128.6 mg of the mixture/capsule, using standard filling equipment.

Claims

1. A controlled release oral pharmaceutical preparation suitable for dosing every 24 hours comprising a substrate comprising a pharmaceutically effective amount of tramadol or a salt thereof and a pharmaceutically effective amount of topiramate; said
5 preparation having a dissolution rate in vitro when measured using the Ph. Eur. Paddle Method at 75 rpm in 900 ml 0.05 M phosphate buffer at 37° C and using HPLC :
between 15 and 35% tramadol released after 1 hour;
between 28 and 48% tramadol released after 2 hours;
10 between 47 and 67% tramadol released after 4 hours;
between 68 and 88% tramadol released after 8 hours;
between 79 and 99% tramadol released after 12 hours;
between 86 and 105% tramadol released after 18 hours;
about 100% tramadol released after 24 hours; by weight,
15 said preparation providing a therapeutic effect for about 24 hours after oral administration.
2. An oral pharmaceutical preparation as in claim 1, said preparation having a dissolution rate as follows :
20 between 20 and 30% tramadol released after 1 hour;
between 33 and 43% tramadol released after 2 hours;
between 52 and 62% tramadol released after 4 hours;
between 73 and 83% tramadol released after 8 hours;
between 84 and 94% tramadol released after 12 hours;
25 between 91 and 100% tramadol released after 18 hours;
about 100% tramadol released after 24 hours; by weight.
3. An oral pharmaceutical preparation as in claim 1, said preparation having a dissolution rate as follows :
30 25% tramadol released after 1 hour;
38% tramadol released after 2 hours;
57% tramadol released after 4 hours;

78% tramadol released after 8 hours;
89% tramadol released after 12 hours;
95,5% tramadol released after 18 hours;
about 100% tramadol released after 24 hours; by weight.

5

4. An oral pharmaceutical preparation according to claims 1-3 wherein the tramadol salt is tramadol hydrochloride.

5. An oral pharmaceutical preparation according to claims 1-4, wherein said substrate is
10 a suitable matrix material in which tramadol or its salt form is incorporated.

6. An oral pharmaceutical preparation according to claim 5 wherein the matrix material is xanthan gum.

15 7. An oral pharmaceutical preparation according to claims 1-6, wherein the preparation comprises two or more phases.

8. An oral pharmaceutical preparation according to claims 1-7 wherein the major part of the tramadol or its salt form and of the topiramate are in different phases of the said
20 pharmaceutical preparation.

9. An oral pharmaceutical preparation according to claims 1-7 wherein the tramadol or its salt form and the topiramate are in different phases of the said pharmaceutical preparation.

25

10. An oral pharmaceutical preparation according to claims 1-9 wherein at least one phase contains the major part of tramadol or a salt thereof and at least another phase contains the major part of topiramate.

30 11. An oral pharmaceutical preparation according to claims 7-10 wherein the phases take the form of layers and at least one layer contains tramadol or a salt thereof and at least another layer contains topiramate.

12. An oral pharmaceutical preparation according to claims 1-11, which has two layers, one layer containing topiramate and another layer containing tramadol or a salt thereof.
- 5 13. An oral pharmaceutical preparation according to claims 1-11 which has three layers, one layer containing topiramate, another layer containing tramadol or a salt thereof, and a third isolator layer in between the tramadol and the topiramate layers.
- 10 14. An oral pharmaceutical preparation according to claims 7-10 wherein at least one of the phases take the form of pellets.
- 15 15. An oral pharmaceutical preparation according to claim 14 wherein all phases take the form of pellets.
16. An oral pharmaceutical preparation according to claims 14 wherein the pellets contain either tramadol or a salt thereof or topiramate.
17. A process for manufacturing an oral pharmaceutical preparation as claimed in claims 1-16 comprising mixing tramadol hydrochloride and topiramate with a suitable
20 substrate.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/14474

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/35 A61K31/135 A61K9/52 A61K9/20 A61K47/36

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal, PAJ, MEDLINE, EMBASE, BIOSIS, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	OBAIDAT A A ET AL: "Controlled release of tramadol hydrochloride from matrices prepared using glyceryl behenate" EUROPEAN JOURNAL OF PHARMACEUTICS AND BIOPHARMACEUTICS, ELSEVIER SCIENCE PUBLISHERS B.V., AMSTERDAM, NL, vol. 52, no. 2, September 2001 (2001-09), pages 231-235, XP004301072 ISSN: 0939-6411 abstract page 232, column 2 -page 234, column 2; figures 1-4	1-17
Y	EP 0 699 436 A (EURO CELTIQUE SA) 6 March 1996 (1996-03-06) cited in the application page 4 -page 5; tables 1-5 -/--	1-17

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

14 April 2004

Date of mailing of the international search report

26/04/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Markopoulos, E

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/14474

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 6 207 197 B1 (ILLUM LISBETH ET AL) 27 March 2001 (2001-03-27) column 4, line 61 -column 5, line 3 column 6, line 26 - line 44 column 9, line 1 - line 28; example 4 ---	1-17
Y	WO 01/13904 A (ORTHO MCNEIL PHARM INC) 1 March 2001 (2001-03-01) cited in the application page 8 -page 14 ---	1-17
Y	US 5 415 871 A (PANKHANIA MAHENDRA G ET AL) 16 May 1995 (1995-05-16) column 2, paragraph 2 column 3, line 53 -column 4, line 45; examples 4,5 ---	1-17
P,X	EP 1 364 649 A (CILAG AG) 26 November 2003 (2003-11-26) page 3, line 4 - line 17 -----	1-4

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/14474

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0699436	A	06-03-1996	DE 4315525 A1 17-11-1994
			GB 2284760 A 21-06-1995
			GB 2287880 A 04-10-1995
			AT 3413 U2 27-03-2000
			AT 138566 T 15-06-1996
			AT 201989 T 15-06-2001
			AU 3995797 A 18-12-1997
			AU 6196394 A 17-11-1994
			CA 2123160 A1 11-11-1994
			CN 1099262 A , B 01-03-1995
			CZ 9401093 A3 16-11-1994
			DE 9422335 U1 13-04-2000
			DE 69400215 D1 04-07-1996
			DE 69400215 T2 31-10-1996
			DE 69427472 D1 19-07-2001
			DE 69427472 T2 08-11-2001
			DE 699436 T1 10-04-1997
			DE 729751 T1 13-03-1997
			DK 624366 T3 01-07-1996
			DK 699436 T3 23-07-2001
			EG 20577 A 30-08-1999
			EP 0624366 A1 17-11-1994
			EP 0699436 A1 06-03-1996
			EP 0729751 A1 04-09-1996
			ES 2088312 T3 01-08-1996
			ES 2159591 T3 16-10-2001
			FI 942092 A 11-11-1994
			FI 20030560 A 14-04-2003
			GR 3020084 T3 31-08-1996
			GR 3036565 T3 31-12-2001
			HU 75703 A2 28-05-1997
			ID 25988 A 10-11-1994
			IL 109460 A 10-03-1998
			IL 119660 A 12-09-2002
			IN 182370 A1 27-03-1999
			IN 182556 A1 01-05-1999
			JP 3045924 B2 29-05-2000
			JP 7053361 A 28-02-1995
			JP 3267561 B2 18-03-2002
			JP 11124327 A 11-05-1999
			JP 3443574 B2 02-09-2003
			JP 2002154954 A 28-05-2002
			NO 941719 A 11-11-1994
			NO 993484 A 11-11-1994
			NO 20013566 A 11-11-1994
			NZ 260408 A 28-05-1996
			PL 176474 B1 30-06-1999
			PT 699436 T 30-10-2001
			SG 67347 A1 21-09-1999
			SK 54194 A3 07-12-1994
US 6207197	B1	27-03-2001	AU 737818 B2 30-08-2001
			AU 7664298 A 11-12-1998
			BR 9809453 A 20-06-2000
			CA 2285580 A1 26-11-1998
			CN 1257425 T 21-06-2000
			EP 0984774 A1 15-03-2000
			WO 9852547 A1 26-11-1998

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/14474

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 6207197	B1		GB 2341550 A ,B	22-03-2000
			HU 0002935 A2	28-12-2000
			JP 2001525848 T	11-12-2001
			NO 995753 A	24-11-1999
			NZ 338105 A	27-04-2001
			PL 336037 A1	05-06-2000
			SK 148999 A3	11-07-2000
			ZA 9804379 A	22-11-1999
WO 0113904	A	01-03-2001	AU 6895800 A	19-03-2001
			BG 106420 A	30-09-2002
			BR 0013439 A	30-04-2002
			CA 2381797 A1	01-03-2001
			CN 1379686 T	13-11-2002
			CZ 20020524 A3	13-11-2002
			EP 1210118 A2	05-06-2002
			HU 0203198 A2	28-02-2003
			JP 2003507421 T	25-02-2003
			NO 20020728 A	05-04-2002
			PL 353266 A1	03-11-2003
			SK 2262002 A3	04-02-2003
			TR 200201067 T2	23-09-2002
			WO 0113904 A2	01-03-2001
			US 6562865 B1	13-05-2003
			ZA 200201403 A	19-05-2003
US 5415871	A	16-05-1995	AT 77548 T	15-07-1992
			AU 608208 B2	28-03-1991
			AU 6762587 A	23-07-1987
			AU 628754 B2	17-09-1992
			AU 7946291 A	12-09-1991
			CA 1313133 C	26-01-1993
			DE 3779933 D1	30-07-1992
			DE 3779933 T2	10-12-1992
			DK 14887 A	19-07-1987
			EP 0234670 A2	02-09-1987
			ES 2042541 T3	16-12-1993
			FI 870155 A ,B,	19-07-1987
			GR 3005796 T3	07-06-1993
			IE 59333 B1	09-02-1994
			IL 81281 A	29-11-1990
			IN 164250 A1	04-02-1989
			JP 1940551 C	09-06-1995
			JP 6069965 B	07-09-1994
			JP 62181227 A	08-08-1987
			KR 9411242 B1	03-12-1994
			MX 4949 A	01-12-1993
			NO 870181 A ,B,	20-07-1987
			NZ 218907 A	29-05-1989
			PH 26408 A	02-07-1992
			PT 84133 A ,B	01-02-1987
			ZA 8700173 A	26-08-1987
EP 1364649	A	26-11-2003	EP 1364649 A1	26-11-2003
			WO 03099273 A1	04-12-2003